

Boca Raton
9980 Central Park Blvd
Suite 202, N
Boca Raton, FL 33428



Provider Order Form

Date: _____

Rituximab (Rituxan, Truxima, Ruxience)

PATIENT INFORMATION

Name:	DOB:
Allergies:	Date of Referral:

ICD-10 code (required): _____ ICD -10 description: _____

NKDA Allergies: _____ Weight lbs/kg: _____

Patient Status: New to Therapy Continuing Therapy Next Due Date (if applicable): _____

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

REFERRAL STATUS

- New Prescription
- Order Renewal
- Does or Frequency Change
- Discontinuation

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Other: _____

PRE-MEDICATION ORDERS

The following are manufacturer recommended premedication regimens:

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- other _____

ADDITIONAL PRE -MEDICATION ORDERS

- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- Other: _____
- Dose: _____ Route: _____
- Frequency: _____

SPECIAL INSTRUCTIONS

THERAPY ADMINISTRATION

Please check preferred product:

- Rituximab(Rituxan) Rituximab-abbs (Truxima)
- Rituximab-pvvr (Ruxience)
- Mix in 0.9% sodium chloride or D5W to final concentration of 1-4mg/ml
 - Dose: 1000mg / _____ mg mg / kg
 - Mix in: 500ml / 250ml other _____
 - Frequency: On Series Day 0 and Series Day 14; repeat series every 24 weeks
 - Other: _____
 - Infusion rate: First infusion in series: 50mg/hr, increasing every 30 minutes by 50mg/hr to maximum of 400mg/hr
 - Subsequent infusion in series: 100mg/hr, increasing every 30 minutes by 100mg/hr to maximum of 400mg
- Flush with 0.9% sodium chloride at the completion of infusion
- Monitor patient for 30 minutes post infusion
- Refills: Zero / for 12 months / _____
(if not indicated order will expire one year from date signed)
- total dosage _____ refill _____

Pre-medicate patients with an antihistamine and acetaminophen prior to dosing. For RA and PV patients, methylprednisolone 100 mg intravenously or its equivalent is recommended 30 minutes prior to each infusion. Screen all patients for HBV infection by measuring HBsAg and anti- Hbc before initiating treatment with RITUXAN. For patients who show evidence of prior hepatitis B infection (HBsAg positive [regardless of antibody status] or HBsAg negative but antiHbc positive), consult with physicians with expertise in managing hepatitis B regarding monitoring and consideration for HBV antiviral therapy before and/or during RITUXAN treatment.

ORDERING PROVIDER

Signature X _____ Date _____

Provider _____ Phone _____ Fax _____