Vermont 28 Park Ave Suite #1A Williston, VT 05495





Idursulfase (Elaprase)

Provider _

Provider Order Form	Date:
PATIENT INFORMATION	
Name:	DOB: SEX: M \square F \square
ICD-10 code (required):	ICD-10 description:
□NKDA Allergies:	Weight lbs/kg:
REFERRAL STATUS	
□New Referral □Referral Renewal □Medication/Order Ch	ange Benefits Verification Only Discontinuation Order
PHYSICIAN	N INFORMATION
Referral Coordinator Name:	Referral Coordinator Email:
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: Fax:
Practice Address:	City: State: Zip Code:
LABORATORY ORDERS	THERAPY ADMINISTRATION
□ CBC □ at each dose □ every □ CMP □ at each dose □ every □ CRP □ at each dose □ every □ Other: □ PRE-MEDICATION ORDERS □ acetaminophen (Tylenol) □ 500mg / □ 650mg / □ 1000mg PO □ cetirizine (Zyrtec) 10mg PO □ loratadine (Claritin) 10mg PO □ diphenhydramine (Benadryl) □ 25mg / □ 50mg □ PO / □ IV □ methylprednisolone (Solu-Medrol) □ 40mg / □ 125mg IV □ hydrocortisone (Solu-Cortef) □ 100mg IV □ Other: □ Dose: Route: Frequency: SPECIAL INSTRUCTIONS SPECIAL INSTRUCTIONS	 Idursulfase (Elaprase) in 100ml 0.9% sodium chloride, intravenous infusion Dose: 0.5mg/kg Route: ☑ intravenous Frequency: once every week The total volume of infusion should be administered over aperiod of 3 hours, which may be gradually reduced to 1 hourif no hypersensitivity reactions are observed. Infuse with a low-protein-binding 0.2 micrometer (OEºm) in-line filter. Flush with 0.9% sodium chloride at infusion completion Patient is required to stay for 30-minute observation period Refills: ☐ Zero / ☐ for 12 months / ☐
NOTES/ADDITIONAL COMMENTS:	
ORDERING PROVIDER	
-,	Б.:
Signature X	Date

Phone Fax _