



(Ultomiris) Ravulizumab-cwvz Infusion orders

Date: _____

PATIENT INFORMATION	
Name:	DOB: SEX: M 🗆 F 🗆
ICD-10 code (required):	ICD-10 description:
□NKDA Allergies:	Weight lbs/kg:
REFERRAL STATUS	
□New Referral □Referral Renewal □Medication/Order Change □Benefits Verification Only □Discontinuation Order	
PHYSICIAN INFORMATION	
Referral Coordinator Name:	Referral Coordinator Email:
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: Fax:
Practice Address:	City: State: Zip Code:
DIAGNOSIS (and ICD 10 code) Myasthenia gravis without (acute) exacerbation ICD-10 Code: G70.00 Myasthenia gravis with (acute) exacerbation ICD-10 Code: G70.01 Other disorders of phosphorus metabolism ICD 10 Code: G70.01 Other disorders of phosphorus metabolism ICD 10 Code: D59.5 Neuromyelitis Optica (NMO), Aquaporin 4 Antibody Positive ICD 10 Code: D59.3 NOTE List Tried & Failed Therapies, including duration of treatment: 1) 2) Immunize patients with meningococcal vaccines at least 2 weeks prior to administering the first dose of ULTOMIRIS, unless the risks of delaying ULTOMIRIS therapy outweigh the risk of developing a meningococcal infection. Comply with the most current National Advisory Committee on Immunization in patients with complement deficiencies.	Ravulizumab-cwvz (Ultomiris) ORDERS Initial Dosing 2,400 mg IV (40k to less than 60kg) 2,700 mg IV (60k to less than 100 kg) 3,000 mg IV (100k or greater kg) Maintenance Dosing 3,000 mg (40k to less than 60kg) IV every 8 weeks starting 2 weeks after initial load 3,300 mg (60k to less than 100 kg) IV every 8 weeks starting 2 weeks after initial load Maintenance Dosing ONLY 3,000 mg (40k to less than 60kg) IV every 8 weeks 3,000 mg (40k to less than 60kg) IV every 8 weeks 3,000 mg (60k to less than 100 kg) IV every 8 weeks 3,000 mg (40k to less than 60kg) IV every 8 weeks 3,000 mg (40k to less than 100 kg) IV every 8 weeks 3,000 mg (40k to less than 100 kg) IV every 8 weeks 3,300 mg (60k to less than 100 kg) IV every 8 weeks 3,300 mg (100k or greater kg) IV every 8 weeks 3,600 mg (100k or greater kg) IV every 8 weeks ADJUST DOSE BASED ON WEIGHT (KG) AT NEXT INFUSION AFTER NOTIFYING DR? ***** Refills*: None □X6 months □X1 year □ Other:*(if not indicated order will expire one year from date signed) REQUIRED DOCUMENTATION: □ This signed order form by the provider □ Patient demographics AND insurance information □ Clinical/ Progress notes supporting primary dx
 Documentation of meningococcal vaccines WITH DATES OF ADMINISTRATION OF MEN B & MEN ACWY OR WITH DATES OF ADMINISTRATION OF MEN ABCWY OR IF NOT FULLY VACCINATED - PHROPHLATIC ANTIBX RX SENT - Is your patient enrolled in the Ultomiris-REMS program?YESNo (if no, must be enrolled to start therapy) Is the ordering PROVIDER enrolleD in the Ultomiris-REMS program?YESNO (if no, must be enrolled to start therapy) = 	
ORDERING PROVIDER	
Signature X	Date
Provider Phone	Fax