Pensacola 41 Fairpoint Drive Suite B Gulf Breeze, FL 32561

Boca Raton 9980 N Central Park Blvd Suite 202 Boca Raton, FL 33428





(Crysvita)

Provider_

Burosumab-twza

Infusion orders	Date:
PATIENT INFORMATION	
Name:	DOB: SEX: M F
ICD-10 code (required):	ICD-10 description:
□NKDA Allergies:	Weight lbs/kg:
REFERRAL STATUS	
□New Referral □Referral Renewal □Medication/Order Ch	
PHYSICIAN INFORMATION	
Referral Coordinator Name:	Referral Coordinator Email:
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: Fax:
Practice Address:	City: State: Zip Code:
DIAGNOSIS (and ICD 10 code) XLH: (familial hypophosphatemia) ICD-10 Code: E83.31 ICD-10 Code: M83.8 ICD-10 Code: M83.8 ICD-10 Code: E83.39 NOTE	Burosumab-twza ORDERS Indication Pediatric XLH (6 months and older) Adult XLH Pediatric TIO 2 years and older Adult TIO *Adult TIO *Adult TIO Medication(check one) Crysvita less than 10 kg
List Tried & Failed Therapies, including duration of treatment: 1) 2) **Referring physician is responsible for monitoring and reviewing the following labs prior to treatment: • Fasting phosphorus level prior to each dose for first 3 doses and administer only if below ULN • Fasting phosphorus level 2-4 weeks after dose modifications If dose adjustments are needed, new order must be sent by provider based on PI dose calculations	Crysvita greater than 10 kg Crysvita Dosing 1 mg/kg SQ rounded to the nearest 1 mg max 90 mg 0.8 mg/kg SQ rounded to the nearest 10 mg max 90 mg 1 mg/kg SQ rounded to the nearest 10 mg max 90 mg 0.4 mg/kg SQ rounded to the nearest 10 mg 2 mg/kg not to exceed 180 mg 0.5 mg/kg not to exceed 180 mg mg/kg (dose may be increased up to 2mg/kg not to exceed 180mg administered every 2weeks) Frequency Every 2 weeks Every 4 weeks Every 4 weeks Every 4 weeks Every 4 weeks Refills*: None □ 36 months □ X1 year □ Other:
REQUIRED DOCUMENTATION: This signed order form by the provider Patient demographics AND insurance information Clinical/Progress notes supporting primary diagnosis Documentation that pt has stopped phos meds and Vit D Fasting serum phosphorus concentration should be below the reference range for age prior to initiation of treatment	
ORDERING PROVIDER Signature X	Date
Signature A	Buic

_____ Phone _____ Fax _