Hackensack 385 Prospect Avenue Suite 101 Hackensack, NJ, 07601

Marlton 127 Church Road Suite 203 Marlton, NJ 08053





Long Branch 422 Morris Avenue Suite 7 Long branch, NJ 07740

Somerset81 Veronica Avenue Suite 202 Somerset NJ 08873

## DIZENAN

BKEWV (eculizumab-aeeb		te:				
	1	<u>INFORMA</u>	ı			
Name:	Phone:		DOB:	SEX: /	M 🗆 F 🗆	
□NKDA Allergies:				Weight lbs/kg:		
	PHYSICIAN	<b>INFORM</b>	ATION			
Physician Name:		Practice Name:				
Address:		Office Contact Name: Office Contact #:				
Phone: Fax:		Email (for updates):				
REFERRAL STATUS						
□New Referral □Referral Renewal □	Medication/Order Ch	ange □Ben	efits Verification	n Only Discont	tinuation Order	
MEDICATION ORDERS  □ gMG who are anti-acetylcholine receptor (ArchR) antibody+ ICD 10:  gMG NEW START dosing (adult dosing)  □ 900 mg weekly for the first 4 weeks, followed by □1,200 mg for the fifth dose 1 week later then □1,200 mg every 2 weeks x • Refills* □None □x6 months □x1year □Other: *(if not indicated order will expire one year from date signed  □ Paroxysmal Nocturnal Hemoglobinuria (PNH) ICD 10: PNH BKEMV NEW START dosing (18 yo and older)		□ atypical Hemolytic Uremic Syndrome (aHUS)  ICD 10:  aHus NEW START dosing (18 yo and older)* □ 900 mg weekly for the first 4 weeks, followed by □ 1,200 mg for the fifth dose 1 week later then □ 1,200 mg every 2 weeks x  • Refills* □None □ x6 months □ x1 year □Other: *(if not indicated order will expire one year from date signed				
						PT wt and dosing
		□ WT	40kg and over 30kg to < 40kg	900mg weekly x 4 doses	1200mg every 2 weeks 900mg at week 3 then	
		□ WT		600mg weekly x 2 doses	900mg every 2 weeks 600mg at week 3 then	
		□ 600 mg weekly for the first 4 weeks, followed by □ 900 mg for the fifth dose 1 week later then □ 900 mg every 2 weeks x • Refills* □None □x6 months □x1year □Other: *(if not indicated order will expire one year from date signed		□ WT	20kg to < 30kg	600mg weekly x 2 doses
□ wt	10kg to < 20kg			600mg weekly x 1 doses	300mg every 2 weeks 300mg at week 2 then	
***Complete or update vaccination from meningococcal bacteria (for serogroups A,C,W,Y, and B) at least 2 weeks prior to the first dose of BKEMV, unless the risks of delaying therapy with BKEMV outweigh the risk of developing a serious infection.Comply with the most current Advisory Committee on Immunization Practices (ACIP) recommendations for vaccinations against meningococcal bacteria in patients receiving a						
				REQUIRED DOCUMENTATION CHECKLIST:		
□ This signed order form by the provider □ Patient demographics AND insurance information □ Clinical/ Progress notes supporting primary dx □ Acetylcholine Receptor Antibody Test Results ( if Myasthenia Gravis)			complement inhibitor. See Warning and Precautions (5.1) for additional guidance on the management of the risk of serious infections caused by meningococcal bacteria.			
		WARNINGS AND PRECAUTIONS https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/761333s001lbl.pdf				
Documentation of meningococcal vaccines  □ WITH DATES OF ADMINISTRATION OF M	1EN B & MEN ACWY <b>O</b>	PR				

- □ WITH DATES OF ADMINISTRATION OF MEN ABCWY **OR**
- □ IF NOT FULLY VACCINATED PROPHYLACTIC ANTIBX RM MUST BE SENT
- $\square$  Is your patient enrolled in the BKEMV REMS program  $\square$  YES  $\square$  NO (If no, must be enrolled to start therapy)
- □ Is the ordering PROVIDER enrolled in the BKEMV REMS program □YES □NO (If no, must be enrolled to start therapy)

## **ORDERING PROVIDER**

Signature X		Date
Provider	Phone	Provider NPI