

Alglucosidase alfa-ngpt (Nexviazyme)

Provider Order Form

Date: _____

PATIENT INFORMATION		
Name:	DOB:	SEX: M ☐ F ☐
ICD-10 code (required):	ICD-10 description:	
☐ NKDA	Allergies:	Weight lbs/kg:

REFERRAL STATUS	
☐ New Referral	☐ Referral Renewal
☐ Medication/Order Change	☐ Benefits Verification Only
☐ Discontinuation Order	

PHYSICIAN INFORMATION	
Referral Coordinator Name:	Referral Coordinator Email:
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: Fax:
Practice Address:	City: State: Zip Code:

LABORATORY ORDERS ☐ CBC ☐ at each dose ☐ every _____ ☐ CMP ☐ at each dose ☐ every _____ ☐ CRP ☐ at each dose ☐ every _____ ☐ Other: _____ PRE-MEDICATION ORDERS ☐ acetaminophen (Tylenol) ☐ 500mg / ☐ 650mg / ☐ 1000mg PO ☐ cetirizine (Zyrtec) 10mg PO ☐ loratadine (Claritin) 10mg PO ☐ diphenhydramine (Benadryl) ☐ 25mg / ☐ 50mg ☐ PO / ☐ IV ☐ methylprednisolone (Solu-Medrol) ☐ 40mg / ☐ 125mg IV ☐ Other: _____ Dose: _____ Route: _____ Frequency: _____ SPECIAL INSTRUCTIONS	THERAPY ADMINISTRATION ☐ Alglucosidase alfa-ngpt (Nexviazyme) in 5% Dextrose, intravenous infusion, final concentration of 0.5 to 4mg/ml, administer with 0.2 micron filter - Dose: ☐ ($\geq$ 30kg) 20mg/kg ☐ ($\leq$ 30kg) 40mg/kg ☐ other _____ - Frequency: every 2 weeks ☐ other _____ - Administer over approximately 4 hours, ☐ other _____ ☐ Flush with 5% Dextrose at the completion of infusion ☐ Patient is required to stay for 30-minute observation period ☐ Patient is NOT required to stay for observation time ☐ Refills: ☐ Zero / ☐ for 12 months / ☐ _____ (if not indicated order will expire one year from date signed) ☐ Total dosages _____ ☐ Refills _____
---	--

NOTES/ADDITIONAL COMMENTS:

ORDERING PROVIDER

Signature X Date _____

Provider _____ Phone _____ Fax _____