

Westerville
575 Copeland Mill Road
Suite# 2F
Westerville, Ohio 43081



Lancaster
2405 Columbus Street
Suite# 210
Lancaster, Ohio 43130

Provider Order Form

Iron (Feraheme/Injectafer/Venofer)

Date: _____

PATIENT INFORMATION

Name:	DOB:
Allergies:	Date of Referral:

ICD-10 code (required): _____ ICD -10 description: _____

☐ NKDA Allergies: _____ Weight lbs/kg: _____

Patient Status: ☐ New to Therapy ☐ Continuing Therapy Next Due Date (if applicable): _____

REFERRAL STATUS: ☐ New Prescription ☐ Order Renewal ☐ Does or Frequency Change ☐ Discontinuation

PROVIDER INFORMATION

Referral Coordinator Name:	Referral Coordinator Email:
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: _____ Fax: _____
Practice Address:	City: _____ State: _____ Zip Code: _____

PREN-MEDICATION ORDERS

- ☐ acetaminophen (Tylenol) ☐ 500mg / ☐ 650mg / ☐ 1000mg PO
- ☐ cetirizine (Zyrtec) 10mg PO
- ☐ loratadine (Claritin) 10mg PO
- ☐ diphenhydramine (Benadryl) ☐ 25mg / ☐ 50mg ☐ PO / ☐ IV
- ☐ methylprednisolone (Solu-Medrol) ☐ 40mg / ☐ 125mg IV
- ☐ Other: _____
- Dose: _____ Route: _____
- Frequency: _____

SPECIAL INSTRUCTIONS

*Closely observe patients for signs and symptoms of hypersensitivity including monitoring of blood pressure and pulse during and after Feraheme administration for at least 30 minutes and until clinically stable following completion of each infusion.

*Observe for signs and symptoms of hypersensitivity during and after Injectafer administration for at least 30 minutes and until clinically stable following completion of each administration. *Monitor patients for signs and symptoms of hypersensitivity during and after Venofer administration for at least 30 minutes and until clinically

THERAPY ADMINISTRATION

- ☐ Ferumoxylol (Feraheme) intravenous infusion
 - Dose & Frequency: ☒ initial 510mg infusion followed by a second 510mg infusion 3-8 days later
 - ☐ Dilute in 50 - 200ml 0.9% sodium chloride or 5% dextrose solution (final concentration 2mg - 8mg per ml)
 - ☐ Infuse over at least 15 minutes
 - ☐ No refills ☐ Other
 - ☐ Ferriccarboxymaltose (Injectafer) intravenous infusion
 - Dose & Frequency:
 - ☐ Patients > 50kg: Two 750mg doses, 7 days apart
 - ☐ Patients < 50kg: Two 15mg/kg doses, 7 days apart
 - ☐ Dilute in no more than 250ml 0.9% sodium chloride
 - ☐ Infuse over at least 15 minutes
 - ☐ No refills ☐ Other
 - ☐ Ironsucrose(Venofer) intravenous infusion
 - Dose:
 - ☐ 100mg in 100ml 0.9% sodium chloride over 30 minutes
 - ☐ 200mg in 100ml 0.9% sodium chloride over 30minutes
 - ☐ 300mg in 250ml 0.9% sodium chloride over 1.5 hours
 - ☐ 400mg in 250ml 0.9% sodium chloride over 2.5 hours
 - ☐ _____
 - Frequency:
 - ☐ Once ☐ Every 2- 3 days x _____ doses
 - ☐ Daily x _____ doses ☐ Weekly x _____ doses
 - ☐ Monthly x _____ doses ☐ Other: _____
- ☒ Flush with 0.9% sodium chloride at the completion of infusion
☒ Patient required to stay for 30 - min observation period
☐ Total doses: ☐ 1 yr ☐ Other

Provider Name (Print) _____ Provider Signature _____ Date _____

ORDERING PROVIDER

Signature X _____ Date _____

Provider _____ Phone _____ Fax _____