

TN
100 Covey Drive
Suite 307
Franklin, TN 37067



Idursulfase (Elaprase) Provider Order Form

Date: _____

PATIENT INFORMATION

Name:	DOB:	SEX: M ☐ F ☐
ICD-10 code (required):	ICD-10 description:	
☐ NKDA Allergies:	Weight lbs/kg:	

REFERRAL STATUS

☐ New Referral ☐ Referral Renewal ☐ Medication/Order Change ☐ Benefits Verification Only ☐ Discontinuation Order

PHYSICIAN INFORMATION

Referral Coordinator Name:	Referral Coordinator Email:
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: Fax:
Practice Address:	City: State: Zip Code:

LABORATORY ORDERS

- ☐ CBC ☐ at each dose ☐ every _____
☐ CMP ☐ at each dose ☐ every _____
☐ CRP ☐ at each dose ☐ every _____
☐ Other: _____

PRE-MEDICATION ORDERS

- ☐ acetaminophen (Tylenol) ☐ 500mg / ☐ 650mg / ☐ 1000mg PO
☐ cetirizine (Zyrtec) 10mg PO
☐ loratadine (Claritin) 10mg PO
☐ diphenhydramine (Benadryl) ☐ 25mg / ☐ 50mg ☐ PO / ☐ IV
☐ methylprednisolone (Solu-Medrol) ☐ 40mg / ☐ 125mg IV
☐ hydrocortisone (Solu-Cortef) ☐ 100mg IV
☐ Other: _____
Dose: _____ Route: _____
Frequency: _____

SPECIAL INSTRUCTIONS

THERAPY ADMINISTRATION

- ☐ **Idursulfase (Elaprase)** in 100ml 0.9% sodium chloride, intravenous infusion
- Dose: 0.5mg/kg
 - Route: ☐ intravenous
 - Frequency: once every week
- The total volume of infusion should be administered over a period of 3 hours, which may be gradually reduced to 1 hour if no hypersensitivity reactions are observed.
- ☐ Infuse with a low-protein-binding 0.2 micrometer (OE^m) in-line filter.
- ☐ Flush with 0.9% sodium chloride at infusion completion
- ☐ Patient is required to stay for 30-minute observation period
- ☐ Refills: ☐ Zero / ☐ for 12 months / ☐ _____
(if not indicated order will expire one year from date signed)
- Total dosages _____
Refills _____

NOTES/ADDITIONAL COMMENTS:

ORDERING PROVIDER

Signature X Date _____

Provider _____ Phone _____ Fax _____