Vermont 28 Park Ave Suite #1A Williston, VT 05495





Date: _____

Canakinumab (Ilaris) Provider Order Form

| Trovider Order rollin | | | |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|----------------|
| PATIENT INFORMATION | | | |
| Name: | DOB: | | SEX: M □ F □ |
| ICD-10 code (required): | ICD-10 description: | | |
| □NKDA Allergies: | | | Weight lbs/kg: |
| REFERRAL STATUS | | | |
| □New Referral □Referral Renewal □Medication/Order Change □Benefits Verification Only □Discontinuation Order | | | |
| PHYSICIAN INFORMATION | | | |
| Referral Coordinator Name: | Referral Coordinator Email: | | |
| Ordering Provider: | Provider NPI: | | |
| Referring Practice Name: | Phone: Fax: | | |
| Practice Address: | City: | State: | Zip Code: |
| OBSERVATION (PLEASE SELECT BELOW) Patient is required to stay for 30 minutes observation period Patient is NOT required to stay for observation time Other: SPECIAL INSTRUCTIONS | THERAPY ADMINISTRATION Canakinumab (Ilaris) For Stills Disease including Adult Onset Stills Disease and Systemic Juvenile Idiopathic Arthritis. 4mg/kg (with a max of 300mg) for patients with a body weight greater than or equal to 7.5kg subcutaneous every 4 weeks Other | | |
| ORDERING PROVIDER Signature X | | Date | |
| 2.0 | | | |
| Provider | Phone | Fax | |