Borough Park 1428 36th Street Suite 107 Brooklyn, NY 11218

Sheepshead Bay 2546 East 17th Street Fl, 1 Brooklyn, NY 11235

Bronx 226 West 238th Street Bronx, NY 10463

Forest Hills 64-05 Yellowstone Blvd CF104 Forest Hills, NY 11375 Long Island City

36-36 33rd Suite 311 Long Island City, NY 11106

New York, NY 10022 FIDI

30 Broad Street Suite 401 New York, NY 10004



Central Park West 115 Central Park West Suite 15 New York, NY 10023





Tarrytown 555 Taxter Road 3rd Floor Elmsford, NY 10523

Port Jefferson 12 Medical Drive Suite B Port Jefferson Station, NY 11776

Staten Island 27 New Dorp Lane Staten Island, NY 10306

Southampton 625 Hampton Road Southampton, NY 11968

1228 E Main Street

Suite A

Riverhead, NY 11901

Holbrook

233 Union Avenue

Suite 207

Holbrook, NY 11741

333 East Shore Road

Suite 201 Manhasset, NY 11030 Rockville Centre

Manhasset

165 North Village Avenue Suite 133 Rockville Center, NY 11570

New Hyde Park 1991 Marcus Ave Suite 110 Lake Success, NY, 11042

7600 Jericho Tpke, Lower Level, Suite C500 Woodbury NY 11797

Massapequa 97 Grand Avenue Gramercy 7 Gramercy Park West Lower Level New York, NY, 10003 Massapequa, NY 11758

PATIENT	ER FORM Date: Vocation was a second with the control of the
Name:	DOB: SEX: M □ F □
Phone:	Preferred Location:
	R INFORMATION
Ordering Provider:	Provider NPI:
Referring Practice Name:	Phone: Fax:
Office Contact:	Address:
mail (required):	Address.
•	RAL STATUS
	Order □Transfer of care – Date of last infusion/Next due date
RECEMBI:	Order - Harister of care - Date of last illiusion/next due date
mpairment (MCI) or mild dementia stage of disease, the population in Please note MRIs to assess for ARIA are required prior to doses 3, 5, 7	7, and 14 and must be sent to Thrivewell with an updated order for the ceed with treatment. Failure to provide the required MRI report at least
■ Diagnosis ICD-10 Check one: □ G31.84 Mild Cognitive impairment, so stated □ G30.0 Alzheimer's Disease with early onset □ G30.1 Alzheimer's Disease with late onset □ G30.8 Other Alzheimer's Disease	PATIENT WEIGHT lbs kg Therapy Administration and Dosing (supplied as 200mg/2mL or 500mg/5mL vial)
☐ G30.9 Alzheimer's Disease, unspecified	All patients will be weighed prior to every infusion and medication will be administered according to that weight, diluted in 0.9%
■ Premeds Select all that apply:	
Premeds Select all that apply: ☐ Acetaminophenmg PO (recommended for first 6 doses)	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses) □ Cetirizine 10mg PO	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses)	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks Doses #5 – 6: 10mg/kg IV every 2 weeks
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses) □ Cetirizine 10mg PO □ Diphenhydramine (check all that apply) 25mg50mgPOIV □ Methylprednisolonemg IV	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks Doses #5 – 6: 10mg/kg IV every 2 weeks Doses #7 – 13: 10mg/kg IV every 2 weeks
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses) □ Cetirizine 10mg PO □ Diphenhydramine (check all that apply) 25mg50mgPOIV □ Methylprednisolonemg IV □ SoluCortefmg IV	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks Doses #5 – 6: 10mg/kg IV every 2 weeks Doses #7 – 13: 10mg/kg IV every 2 weeks Doses #7 – 13: 10mg/kg IV every 2 weeks Doses #: 10mg/kg IV every 2 weeks
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses) □ Cetirizine 10mg PO □ Diphenhydramine (check all that apply) 25mg50mgPOIV □ Methylprednisolonemg IV	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks Doses #5 – 6: 10mg/kg IV every 2 weeks Doses #7 – 13: 10mg/kg IV every 2 weeks
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses) □ Cetirizine 10mg PO □ Diphenhydramine (check all that apply) 25mg50mgPOIV □ Methylprednisolonemg IV □ SoluCortefmg IV	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks Doses #5 – 6: 10mg/kg IV every 2 weeks Doses #7 – 13: 10mg/kg IV every 2 weeks Doses #: 10mg/kg IV every 2 weeks MAINTENANCE DOSING (CAN ONLY BE SELECTED ONCE PATIENT HAS COMPLETED 18 MONTHS OF STANDARD DOSING AT THE
■ Premeds Select all that apply: □ Acetaminophenmg PO (recommended for first 6 doses) □ Cetirizine 10mg PO □ Diphenhydramine (check all that apply) 25mg50mgPOIV □ Methylprednisolonemg IV □ SoluCortefmg IV	will be administered according to that weight, diluted in 0.9% NS IVPB over 1 hour. ONLY ONE DOSE SET CAN BE SELECTED AND A NEW ORDER WILL BE REQUIRED FOR EACH SUBSEQUENT SET Doses #1 – 2: 10mg/kg IV every 2 weeks Doses #3 – 4: 10mg/kg IV every 2 weeks Doses #5 – 6: 10mg/kg IV every 2 weeks Doses #7 – 13: 10mg/kg IV every 2 weeks Doses #: 10mg/kg IV every 2 weeks MAINTENANCE DOSING (CAN ONLY BE SELECTED ONCE PATIENT HAS COMPLETED 18 MONTHS OF STANDARD DOSING AT THE 2 WEEK INTERVAL PER THE PI) Doses #: 10mg/kg IV every 4 weeks

• All patients will be kept for 30 minutes of monitoring following completion of the first infusion and then on a case-by-case basis subsequently for any clinical

Additional Orders: By signing this order, you agree to the following orders unless otherwise noted.

- · Hold infusion and notify provider if patient reports: Headache, dizziness, nausea, vision changes, new or worsening confusion, balance concerns, or change in mentation.
- Infusion/allergic reactions may be managed by clinical staff in real time per facility protocol. The prescriber's office will be notified in real time of any infusion reactions

Provider Signature:	Date:
Notes/Additional Comments:	